

SU RUZENA 2-row

High yield, short, earlier ripening



Advantages:

- Yield and high allround-use as feed barley in all regions
- Early heading and long grain filling are advantages for trustable yield
- Outstanding lodging tolerance and disease resistance

Cultivation:

Very interesting for farms with organic fertilization due to good standing power

Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

Plant Development



Vitality



Plant Health



Quality



SU RUZENA 2-row

High yield, short, earlier ripening

Plant development and vitality:

Ear emergence	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Maturity	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Yield structure:

Ears/m ²	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Grains/ear	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
TGW	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Sowing:

Time of sowing	20th of September until 25th of Octobre
Seed rate	locally adapted

Straw:

Plant height	(5) medium	
Lodging resistance		<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Necking resistance		<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Plant growth regulators	lower	

Disease resistance:

Mildew	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Net blotch	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Rhynchosporium	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Plant protection	Early sowing: insecticide against aphids (BYDV), splitted application in EC 32 + EC 49/59								

Quality:

Grading (> 2,2 mm)	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Grading (> 2,5 mm)	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Hectolitre weight	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Raw protein content	(2) low to very low