

YVONNE 2-row

Feed barley with best health and lodging tolerance



Advantages:

- Maritime variety type with high yield untreated and treated
- High above average in all terms all kind of disease tolerances, especially good in brown rust
- Best lodging tolerance (no ear break, no stem break)

Cultivation:

Grow on locations with slow ripening conditions
No sowing in conditions with strong frost (without expected snow cover)
Avoid too early sowing in regions with high frost risk

Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

Plant Development

Maturity 

Vitality

Lodging resistance 

Drought resistance 

Necking resistance 

Plant Health

Mildew 

Brown rust 

Quality

Hectolitre weight 

Grading (> 2,5 mm) 

YVONNE 2-row

Feed barley with best health and lodging tolerance

Plant development and vitality:

Ear emergence	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Maturity	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Winter hardiness	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Drought resistance	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Yield structure:

Ears/m ²	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Grains/ear	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
TGW	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Sowing:

Time of sowing	25th of September until 25th of Oct locally adapted
Seed rate	locally adapted

Straw:

Plant height	(4) short to medium
Lodging resistance	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Necking resistance	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Plant growth regulators	lower

Disease resistance:

Mildew	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Net blotch	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Rhynchosporium	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>
Brown rust	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>	<div></div>

Quality:

Grading (> 2,2 mm)	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Grading (> 2,5 mm)	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Hectolitre weight	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Raw protein content	(2) low to very low