

# HYBERY<sup>B</sup>

Recommended by farmers: Healthy and productive



## Advantages:

- High yielding variety suitable for extensive or intensive technology
- Outstanding resistance to all root, leaf and ear diseases
- Resistance to Eye Spot due to Pch1 gene
- Resistance to Orange blossom midge
- Superior baking quality
- Tolerant to CTU

## Cultivation:

High potential in grain and straw yield for all locations  
Recommended for wheat after wheat crop rotations

## Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

### Plant Development



### Vitality

Winter hardiness high to very high

Drought resistance high



### Plant Health



### Quality



# HYBERY<sub>B</sub>

Recommended by farmers: Healthy and productive

## Plant development and vitality:

Ear emergence	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Maturity	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Winter hardiness	high to very high
Drought resistance	high
CTU-tolerant	Tolerant

## Yield structure:

Yield structure	Compensation type
Ears/m <sup>2</sup>	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Grains/ear	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
TGW	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

## Sowing:

Time of sowing	very early to optimum
Seed rate	100 -150 plants/m <sup>2</sup>

## Straw:

Plant height	6 (medium to tall)
Lodging resistance	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Plant growth regulator	locally adapted

## Disease resistance:

Mildew	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Septoria leaf blotch	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Yellow rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Brown rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
DTR	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Fusarium	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Eye spot	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

# HYBERY<sup>B</sup>

Recommended by farmers: Healthy and productive

### Fungicides for intensive cultivation:

Early treatment	only occasionally required
Final treatment	according standard

### Quality:

Hagberg / Falling number	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Hagberg stability	Good
Zeleny / Sedimentation value	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Protein content	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Water absorption	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Flour extraction	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Loaf volume	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>