

# HYPOCAMP<sup>B</sup>

Healthy hybrid, low input with biscuit quality



## Advantages:

- Outstanding resistance to all leaf- and ear diseases
- Good specific weight - soft milling wheat
- Good winter hardiness

## Cultivation:

Use appropriated nitrogen input for biscuit use

## Short profile:

---- = very low resistance/early/short,

++++ = very high resistance/late/long

### Plant Development

Maturity



### Vitality

Winter hardiness

medium to high

Drought resistance

high

Lodging resistance



### Plant Health

Fusarium



### Quality

Hagberg / Falling number



Protein content



# HYPOCAMP<sup>B</sup>

Healthy hybrid, low input with biscuit quality

## Plant development and vitality:

Ear emergence	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Maturity	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Winter hardiness	medium to high
Drought resistance	high

## Yield structure:

Ears/m <sup>2</sup>	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Grains/ear	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
TGW	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

## Sowing:

Time of sowing	early to optimum
Seed rate	130-150

## Straw:

Lodging resistance	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Plant growth regulator	locally adapted, slightly more

## Disease resistance:

Mildew	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Septoria leaf blotch	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Yellow rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Brown rust	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
DTR	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Fusarium	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Eye spot	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>

## Fungicides for intensive cultivation:

Early treatment	only occasionally required
Final treatment	standard

# HYPOCAMP<sup>B</sup>

Healthy hybrid, low input with biscuit quality

## Quality:

Hagberg / Falling number	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Hagberg stability	good
Zeleny / Sedimentation value	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Protein content	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Water absorption	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Flour extraction	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>
Loaf volume	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>